

## Problem 1

Reproduce the formatting of the following equation. You might want to use the following commands:

`\align;` `\intertext;` `\left;` `\right;` `\sum;` `\prod` `\dots;` `\frac;` `\log;`

$$\frac{\sum_{i=1}^n \log(a_i)}{n} = \frac{\log(a_1) + \log(a_2) + \dots + \log(a_n)}{n} \tag{1}$$

$$= \frac{\log(a_1 a_2 \dots a_n)}{n} \tag{2}$$

$$= \log \left[ \left( \prod_{i=1}^n a_i \right)^{\frac{1}{n}} \right] \tag{3}$$

because  $\log(MN) = \log(M) + \log(N)$  and  $\log(M^k) = k \log(M)$

$$= \log \left( \prod_{i=1}^n \sqrt[n]{a_i} \right) \tag{4}$$

## Problem 2

Suppose a crime has been committed. Blood is found at the scene for which there is no innocent explanation. It is of a type which is present in 1% of the population.

1. The prosecutor claims: “There is a 1% chance that the defendant would have the crime blood type if he were innocent. Thus there is a 99% chance that he guilty”. This is known as the prosecutor’s fallacy. What is wrong with this argument?
2. The defender claims: “The crime occurred in a city of 800,000 people. The blood type would be found in approximately 8000 people. The evidence has provided a probability of just 1 in 8000 that the defendant is guilty, and thus has absolutely no bearing on the investigation.” This is known as the defender’s fallacy. What is wrong with this argument?

## Problem 3

My neighbor has two children. Assume that the gender of a child is a coin flip. Let the genders of the children be  $G_1$  and  $G_2$ . For both questions, write the probability symbolically (e.g. “ $P(A|B)$ ”) and give the value.

## Assignment 2

---

1. Suppose I happen to see one of his children run by, and it is a boy. What is the probability that the other child is a girl?
2. Suppose instead that I ask him whether he has any boys, and he says yes. What is the probability that one child is a girl?

### Problem 4

Consider the Numbers game with one-sided interval hypotheses  $h_{\leq x}$  for numbers 1 up to 10, where  $h_{\leq x} = \{1, 2, \dots, x\}$ ,  $x \in \{1 \dots 10\}$ . Assume we have a uniform prior over  $h$ .

1. Show that the  $h_{mle} = h_{\leq \max(S)}$  for a given set of numbers  $S = \{x_1, x_2, \dots\}$ .
2. Briefly say why the MAP estimate = MLE.
3. For Parts 3 and 4, let's suppose  $S = \{5, 9\}$ . What is the plug-in approximation of the posterior predictive distribution for new data point  $x$ ?
4. What is the full posterior predictive distribution?